

CHAPTER – 25

CALORIMETRY

1. Mass of aluminium = 0.5kg,
 Mass of Iron = 0.2 kg
 Sp heat of Iron = $100^{\circ}\text{C} = 373^{\circ}\text{k}$.
 Sp heat of Iron = 470J/kg-k
 Heat again = $0.5 \times 910(T - 293) + 0.2 \times 4200 \times (343 - T)$
 $= (T - 292)(0.5 \times 910 + 0.2 \times 4200)$ Heat lost = $0.2 \times 470 \times (373 - T)$
 \therefore Heat gain = Heat lost
 $\Rightarrow (T - 292)(0.5 \times 910 + 0.2 \times 4200) = 0.2 \times 470 \times (373 - T)$
 $\Rightarrow (T - 293)(455 + 8400) = 49(373 - T)$
 $\Rightarrow (T - 293)\left(\frac{1295}{94}\right) = (373 - T)$
 $\Rightarrow (T - 293) \times 14 = 373 - T$
 $\Rightarrow T = \frac{4475}{15} = 298 \text{ k}$
 $\therefore T = 298 - 273 = 25^{\circ}\text{C}$. The final temp = 25°C .
2. mass of Iron = 100g
 mass of water = 240g
 $S_{\text{iron}} = 470\text{J/kg}^{\circ}\text{C}$
 So, $\frac{100}{1000} \times 470 \times (\theta - 60) = \frac{250}{1000} \times 4200 \times (60 - 20)$
 $\Rightarrow 470 - 47 \times 60 = 25 \times 42 \times 40$
 $\Rightarrow \theta = 4200 + \frac{2820}{47} = \frac{44820}{47} = 953.61^{\circ}\text{C}$
3. The temp. of A = 12°C The temp. of B = 19°C
 The temp. of C = 28°C The temp of $\Rightarrow A + B = 16^{\circ}$
 The temp. of $\Rightarrow B + C = 23^{\circ}$
 In accordance with the principle of calorimetry when A & B are mixed
 $M_{\text{CA}}(16 - 12) = M_{\text{CB}}(19 - 16) \Rightarrow CA4 = CB3 \Rightarrow CA = \frac{3}{4}CB \quad \dots(1)$
 And when B & C are mixed
 $M_{\text{CB}}(23 - 19) = M_{\text{CC}}(28 - 23) \Rightarrow 4CB = 5CC \Rightarrow CC = \frac{4}{5}CB \quad \dots(2)$
 When A & c are mixed, if T is the common temperature of mixture
 $M_{\text{CA}}(T - 12) = M_{\text{CC}}(28 - T)$
 $\Rightarrow \left(\frac{3}{4}\right)CB(T - 12) = \left(\frac{4}{5}\right)CB(28 - T)$
 $\Rightarrow 15T - 180 = 448 - 16T$
 $\Rightarrow T = \frac{628}{31} = 20.258^{\circ}\text{C} = 20.3^{\circ}\text{C}$

